# SMURF2 Antibody

Catalog No: #37124

Package Size: #37124-1 50ul #37124-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


## Description

| Product Name          | SMURF2 Antibody  |
|-----------------------|--|
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Purification          | Antigen affinity purification.   |
| Applications          | IHC  |
| Species Reactivity    | Hu Ms  |
| Specificity           | The antibody detects endogenous levels of total SMURF2 protein.                                      |
| Immunogen Type        | Peptide  |
| Immunogen Description | Synthetic peptide corresponding to a region derived from internal residues of human SMAD specific E3 |
|                       | ubiquitin protein ligase 2   |
| Target Name           | SMURF2   |
| Other Names           | DKFZp686F0270;SMUF2; SMURF2  |
| Accession No.         | Swiss-Prot#: Q9HAU4NCBI Gene ID: 64750Gene Accssion: NP_073576                                       |
| Concentration         | 1.7mg/ml   |
| Formulation           | Rabbit IgG in pH7.3 PBS, 0.05% NaN3, 50% Glycerol.   |
| Storage               | Store at -20°C   |


### **Application Details**

Immunohistochemistry: 1:25-1:100

#### **Images**



Immunohistochemical analysis of paraffin-embedded Human gastric cancer tissue using #37124 at dilution 1/15.



Immunohistochemical analysis of paraffin-embedded Human liver cancer tissue using #37124 at dilution 1/15.

#### Background

E3 ubiquitin-protein ligase SMURF2?is an?enzyme?that in humans is encoded by the?SMURF2?gene. E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Interacts with SMAD1 and SMAD7 in order to trigger their ubiquitination and proteasome-dependent degradation. In addition, interaction with SMAD7 activates autocatalytic degradation, which is prevented by interaction with SCYE1. Forms a stable complex with the TGF-beta receptor-mediated phosphorylated SMAD2 and SMAD3.

#### **Published Papers**

el at., Everolimus Alleviates Renal Allograft Interstitial Fibrosis by Inhibiting Epithelial-to-Mesenchymal Transition Not Only via Inducing Autophagy but Also via Stabilizing IkB- $\alpha$ .In Front Immunol. 2022 Jan 24 by Zeping Gui , Chuanjian Suo, et al. PMID: 35140705, , (2022)

PMID:35140705

Note: This product is for in vitro research use only and is not intended for use in humans or animals.