eNOS (Phospho-Ser615) Antibody

Catalog No: #12137

Package Size: #12137-1 50ul #12137-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | |
|-----------------------|---|
| Product Name | eNOS (Phospho-Ser615) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous levels of eNOS only when phosphorylated at serine 615. |
| Immunogen Type | peptide |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 615 (F-N-S(p)-I-S) derived from Human eNOS. |
| Target Name | eNOS |
| Modification | Phospho |
| Other Names | Constitutive NOS; EC 1.14.13.39; EC-NOS; ECNOS; Endothelial NOS; NOS; type III; NOS3; NOSIII; |
| | Nitric-oxide synthase; endothelial; cNOS |
| Accession No. | Swiss-Prot#:P29474;NCBI Gene#:4846 |
| SDS-PAGE MW | 140kd |
| Concentration | 1.0mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide |
| | and 50% glycerol. |
| Storage | Store at -20°C |

Application Details

Western blotting: 1:500~1:3000

Images



Western blot analysis of extracts from K562 cells, treated with EGF (40nM, 30mins), using eNOS (Phospho-Ser615) antibody #12137. The lane on the right is treated with the synthesized peptide.

Background

Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets.

Isoform eNOS13C: Lacks eNOS activity, dominant-negative form that may down-regulate eNOS activity by forming heterodimers with isoform 1.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.