p16-INK4a (Phospho-Ser152) Antibody

Catalog No: #12148

Package Size: #12148-1 50ul #12148-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| Description | |
|-----------------------|---|
| Product Name | p16-INK4a (Phospho-Ser152) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB IHC |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of p16-INK4a only when phosphorylated at serine 152. |
| Immunogen Type | peptide |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 152 (G-P-S(p)-D-I) derived from Human p16-INK4a. |
| Target Name | p16-INK4a |
| Modification | Phospho |
| Other Names | CD2A1; CDK4I; CDKN2; CDKN2A; CDN2; Cyclin-dependent kinase 4 inhibitor A; cyclin-dependent kinase |
| | inhibitor 2A; MTS1; Multiple tumor suppressor 1; p14ARF; p16(INK4a); p16-INK4; P16INK4A |
| Accession No. | Swiss-Prot#:P42771;NCBI Gene#:1029 |
| SDS-PAGE MW | 20kd |
| Concentration | 1.0mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide |
| | and 50% glycerol. |

Application Details

Western blotting: 1:500~1:3000
Immunohistochemistry: 1:50~1:100


Images

Storage



Western blot analysis of extracts from HeLa cells, treated with EPO (20U/ml, 15mins), using p16-INK4a (Phospho-Ser152) antibody #12148. The lane on the right is treated with the synthesized peptide.

Store at -20°C



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using p16-INK4a (Phospho-Ser152) antibody #12148. The picture on the right is treated with the synthesized peptide.

Background

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.