

Angiotensin-2 Antibody

Catalog No: #48274



Package Size: #48274-1 50ul #48274-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Angiotensin-2 Antibody
Host Species	Mouse
Clone No.	4A2
Purification	ProA affinity purified
Applications	WB, IP, IF, IHC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	peptide
Other Names	AGPT 2 antibody Agpt2 antibody ANG 2 antibody ANG-2 antibody ANG2 antibody Angiotensin 2a antibody Angiotensin 2B antibody Angiotensin-2 antibody Angiotensin2 antibody ANGP2_HUMAN antibody ANGPT 2 antibody Angpt2 antibody Tie2 ligand antibody
Accession No.	Swiss-Prot#:O15123
Calculated MW	62-70kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at 4°C


Application Details

WB: 1:100-1:1,000


IHC: 1:50-500

IP: 1-2 µg per 100-500 µg of total protein(1 ml of cell lysate)

Images



Western Blot analysis of Ang-2 expression in HUV-EC-C (A) and TF-1 (B) whole cell lysates.



Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing cytoplasmic staining of trophoblastic cells at low (A) and high (B) magnification.

Background

Tie-1 and Tie-2 (also designated Tek) are novel cell surface receptor tyrosine kinases. The extracellular domain of Tie-1 has an unusual multidomain structure consisting of a cluster of three epidermal growth factor homology motifs localized between two immunoglobulin-like loops, which are followed by three Fibronectin type III repeats next to the transmembrane region. Angiopoietin-1 (Ang-1) is a secreted ligand for Tie-2. Preliminary biochemical analyses of Ang-1 reveal a potential Fibrinogen-like domain at the carboxy-terminus and coiled-coil regions in the amino-terminus. Ang-1 is an angiogenic factor that is thought to be involved in endothelial development. A related protein, angiopoietin-2 (Ang-2), has been identified as a naturally occurring antagonist of Ang-1 activation of Tie-2. In adult tissue, Ang-2 expression seems to be restricted to sites of vascular remodeling.

References

1. Partanen, J., et al. 1992. A novel endothelial cell surface receptor tyrosine kinase with extracellular epidermal growth factor homology domains. *Mol. Cell. Biol.* 12: 1698-1707.
2. Dumont, D.J., et al. 1992. Tek, a novel tyrosine kinase gene located on mouse chromosome 4, is expressed in endothelial cells and their presumptive precursors. *Oncogene* 7: 1471-1480.
3. Dumont, D.J., et al. 1993. The endothelial-specific receptor tyrosine kinase, Tek, is a member of a new subfamily of receptors. *Oncogene* 8: 1293-1301.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.