# Phospho-Tau(T231) Rabbit mAb

Catalog No: #13381

Package Size: #13381-1 50ul #13381-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

# Description


Product Name	Phospho-Tau(T231) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	SC58-08
Purification	ProA affinity purified
Applications	WB, IHC,IP
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Thr231 of human Tau.
Other Names	Al413597 antibody AW045860 antibody DDPAC antibody FLJ31424 antibody FTDP 17 antibody G protein
	beta1/gamma2 subunit interacting factor 1 antibody MAPT antibody MAPTL antibody MGC134287 antibody
	MGC138549 antibody MGC156663 antibody Microtubule associated protein tau antibody Microtubule
	associated protein tau isoform 4 antibody Microtubule-associated protein tau antibody MSTD antibody Mtapt
	antibody MTBT1 antibody MTBT2 antibody Neurofibrillary tangle protein antibody Paired helical filament tau
	antibody Paired helical filament-tau antibody PHF tau antibody PHF-tau antibody PPND antibody
	PPP1R103 antibody Protein phosphatase 1, regulatory subunit 103 antibody pTau antibody RNPTAU
	antibody TAU antibody TAU_HUMAN antibody Tauopathy and respiratory failure, included antibody
Accession No.	Swiss-Prot#:P10636
Calculated MW	46 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

## **Application Details**

WB: 1:1,000

IHC:1:50-1:200

## **Images**



Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti- phospho -Tau(T231) antibody. Counter stained with hematoxylin.

#### Background

Tau, also known as MAPT (microtubule-associated protein tau), MAPTL, MTBT1 or TAU, is a 758 amino acid protein that localizes to the cytoplasm, as well as to the cytoskeleton and the cell membrane, and contains four Tau/MAP repeats. Expressed in neuronal tissue and existing as multiple alternatively spliced isoforms, Tau functions to promote microtubule assembly and stability and is thought to be involved in the maintenance of neuronal polarity. Tau may also link microtubules with neural plasma membrane components and, addition to its role in microtubule stability, is also necessary for cytoskeletal plasticity. Tau is highly subject to a variety of post-translational modifications, including phosphorylation on serine and threonine residues, polyubiquitination (and subsequent proteasomal degradation) and glycation of specific Tau isoforms. Defects in the gene encoding Tau are associated with Alzheimers disease, pallido-ponto-nigral degeneration (PPND), corticobasal degeneration (CBD) and progressive supranuclear palsy (PSP).

## **References**

- 1. Wang, HY. et al. 2012. Reducing amyloid-related Alzheimer's disease pathogenesis by a small molecule targeting filamin A. J. Neurosci. 32: 9773-9784.
- 2. Kamnaksh, A. et al. 2012. Neurobehavioral, cellular, and molecular consequences of single and multiple mild blast exposure. Electrophoresis. 33: 3680-3692.

#### **Published Papers**

el at., Physiological clearance of  $A\beta$  by spleen and splenectomy aggravates Alzheimer-type pathogenesis. In Aging Cell on 2022 Jan by Zhong-Yuan Yu, Dong-Wan Chen, et al.. PMID:34939734, , (2022)

PMID:34939734

el at., Physiological clearance of  $A\epsilon^{\circ}$ Y by spleen and splenectomy aggravates Alzheimer-type pathogenesis. In Aging Cell on 2021 Dec 23 by Zhong-Yuan Yu, Dong-Wan Chen, et al.. PMID:34939734, , (2021)

PMID:34939734

el at., 14-3-3 $\epsilon$   $^3$  Captures SET in the Cytoplasm, Mediating Tau Pathology and Cognitive Impairments., , (2021)

PMID:

Note: This product is for in vitro research use only and is not intended for use in humans or animals.