Product Datasheet

Vesicle-associated membrane protein 7 Polyclonal Antibody

Catalog No: #42365




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Vesicle-associated membrane protein 7 Polyclonal Antibody
Rabbit
Polyclonal
Caprylic Acid Ammonium Sulfate Precipitation purified
IHC
Hu
The antibody detects endogenous level of total Vesicle-associated membrane protein 7 polyclonal antibody.
protein
Recombinant human Vesicle-associated membrane protein 7 protein
Vesicle-associated membrane protein 7
Synaptobrevin-like protein 1, Tetanus-insensitive VAMP
Swiss-Prot#: P51809
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Store at -20°C

Application Details

Immunohistochemistry: 1:20 - 1:200

Images



Immunohistochemical analysis of paraffin-embeded human colorectal carcinoma using #42365 at dilution of 1:100.

Background

Involved in the targeting and/or fusion of transport vesicles to their target membrane during transport of proteins from the early endosome to the lysosome. Required for heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion. Required for calcium regulated lysosomal exocytosis. Involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. Required for exocytosis of mediators during eosinophil and neutrophil degranulation, and target cell killing by natural killer cells. Required for focal exocytosis of late endocytic vesicles during phagosome formation.

[1]"A synaptobrevin-like gene in the Xq28 pseudoautosomal region undergoes X inactivation."D'Esposito M., Ciccodicola A., Gianfrancesco F., Esposito T., Flagiello L., Mazzarella R., Schlessinger D., D'Urso M.Nat. Genet. 13:227-229(1996)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.