

# Fyn (Phospho-Tyr530) Conjugated Antibody

Catalog No: #C12008

Package Size: #C12008-AF350 100ul #C12008-AF405 100ul #C12008-AF488 100ul #C12008-AF555 100ul #C12008-AF594 100ul #C12008-AF647 100ul #C12008-AF680 100ul #C12008-AF750 100ul #C12008-Biotin 100ul #C12008-Conjugated 50ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Fyn (Phospho-Tyr530) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB, IH, IF
Species Reactivity	Human, Mouse, Rat, Bovine, Pig, Zebrafish
Specificity	The antibody detects endogenous level of Fyn only when phosphorylated at Tyrosine 530.
Immunogen Description	Peptide sequence around phosphorylation site of Tyrosine 530 (P-Q-Y(p)-Q-P) derived from Human Fyn.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	SLK;SYN;p59-FYN
Accession No.	Swiss-Prot#:P06241 NCBI Gene ID:2534 NCBI mRNA#:NM_002037.5? NCBI Protein#:NP_002028.1
Calculated MW	59
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

WB: 1:50-1:200

IF: 1:50-1:200

## Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

## Background

Non-receptor tyrosine-protein kinase that plays a role in many biological processes including regulation of cell growth and survival, cell adhesion, integrin-mediated signaling, cytoskeletal remodeling, cell motility, immune response and axon guidance. Inactive FYN is phosphorylated on its C-terminal tail within the catalytic domain. Following activation by PKA, the protein subsequently associates with PTK2/FAK1, allowing PTK2/FAK1 phosphorylation, activation and targeting to focal adhesions. Involved in the regulation of cell adhesion and motility through phosphorylation of CTNNB1 (beta-catenin) and CTNND1 (delta-catenin). Regulates cytoskeletal remodeling by phosphorylating several proteins including the actin regulator WAS and the microtubule-associated proteins MAP2 and MAPT. Promotes cell survival by phosphorylating AGAP2/PIKE-A and preventing its apoptotic cleavage. Participates in signal transduction pathways that regulate the integrity of the glomerular slit diaphragm (an essential part of the glomerular filter of the kidney) by phosphorylating several slit diaphragm components including NPHS1, KIRREL and TRPC6. Plays a role in neural processes by phosphorylating DPYSL2, a multifunctional adapter protein within the central nervous system, ARHGAP32, a regulator for Rho family GTPases implicated in various neural functions, and SNCA, a small pre-synaptic protein. Participates in the downstream signaling pathways that lead to T-cell differentiation and proliferation following T-cell receptor (TCR) stimulation. Also participates in negative feedback regulation of TCR signaling

through phosphorylation of PAG1, thereby promoting interaction between PAG1 and CSK and recruitment of CSK to lipid rafts. CSK maintains LCK and FYN in an inactive form. Promotes CD28-induced phosphorylation of VAV1.

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.