

## YBOX1 / YB-1 Conjugated Antibody

Catalog No: #C48232



Package Size: #C48232-AF350 100ul #C48232-AF405 100ul #C48232-AF488 100ul #C48232-AF555 100ul #C48232-AF594 100ul  
 #C48232-AF647 100ul #C48232-AF680 100ul #C48232-AF750 100ul #C48232-Biotin 100ul #C48232-Conjugated 50ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	YBOX1 / YB-1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB, IF, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	peptide
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	BP 8 antibody CBF-A antibody CCAAT binding transcription factor I subunit A antibody CCAAT-binding transcription factor I subunit A antibody CSDA2 antibody CSDB antibody DBPB antibody DNA binding protein B antibody DNA-binding protein B antibody EFl-A antibody Enhancer factor I subunit A antibody MDR NF1 antibody MGC104858 antibody MGC110976 antibody MGC117250 antibody NSEP 1 antibody NSEP1 antibody Nuclease sensitive element binding protein 1 antibody Nuclease-sensitive element-binding protein 1 antibody p50 antibody Q15905 antibody Y-box binding protein 1 antibody Y-box transcription factor antibody Y-box-binding protein 1 antibody YB 1 antibody YB-1 antibody YBOX1_HUMAN antibody YBX 1 antibody ybx1 antibody
Accession No.	Swiss-Prot#:Q9Y2T7
Calculated MW	40kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

WB: 1:50-1:200

IF: 1:50-1:200

FC: 1:50-1:200

## Background

MSY2 and YB-2 (MSY3,4) belong to the Y-box family of multifunctional proteins that regulate both transcription and translation. Y-box proteins interact with a wide variety of nucleic acid structures to act as transcription factors and mRNA masking proteins. The modular structure of Y-box proteins includes a highly conserved N-terminal cold-shock domain (CSD, equivalent to the bacterial cold-shock proteins) and four basic C-terminal domains containing arginine clusters and aromatic residues. MSY2 is expressed in testis and ovary where it may repress translation of parental mRNA. The gene encoding human MSY2 maps to chromosome 17p11.2-13.1. YB-2 (MSY3,4 in mouse) is also known as DNA binding protein A and is highly expressed in the testis, heart and muscle. MSY2 and YB-2 bind to the consensus sequence 5'-UCCAUCA-3' contained in the Y-box element.

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.