

IKK γ Polyclonal Antibody

Catalog No: #46773



Package Size: #46773-1 50ul #46773-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	IKK γ Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous IKK γ
Immunogen Type	protein
Immunogen Description	Synthetic peptide from human protein at AA range: 60-120
Target Name	IKK γ
Other Names	NF-kappa-B essential modulator (NEMO) (FIP-3) (I κ B kinase-associated protein 1) (IKKAP1) (Inhibitor of nuclear factor kappa-B kinase subunit gamma) (I-kappa-B kinase subunit gamma) (IKK-gamma) (IKKG) (I κ B kinase subunit gamma) (NF-kappa-B essential modifier)
Accession No.	Swiss-Prot:Q9Y6K9NCBI Gene ID:8517
Calculated MW	50 kDa
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C

Application Details


Western blotting: 1B \times B1500-1B \times B12000

Immunohistochemistry: IHC 1:50-300


Images




Western blot analysis of mouse-kidney mouse-brain mouse-lung mouse-heart 293T lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200

Note: This product is for in vitro research use only and is not intended for use in humans or animals.