DR3 Antibody

Catalog No: #24032



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escr	מוי	tion
\boldsymbol{L}	COUL	ıv	เเบเ

Product Name	DR3 Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	DEAE purified	
Applications	ELISA WB	
Species Reactivity	Hu Ms	
Specificity	DR3 has no cross reaction to other death receptors.	
Immunogen Type	Peptide	
Immunogen Description	Raised against a peptide corresponding to amino acids 25 to 40 of human DR3 precusor.	
Target Name	DR3	
Other Names	Wsl-1, Apo-3	
Accession No.	AAQ88676	
Concentration	1mg/ml	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	


Application Details

Predicted MW: 59 kd


Images



Western blot analysis of DR3 in Jurkat total cell lysate with DR3 antibody at 1:1000 dilution.



Immunofluorescence of DR3 in Jurkat cells with DR3 antibody at 5 μ g/ml.



Immunocytochemistry of DR3 in Jurkat cells with DR3 antibody at 5 $\mu g/ml$.

Background

Apoptosis, or programmed cell death, occurs during normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by certain cytokines including TNF and Fas ligand of the TNF family through their death domain containing receptors, TNFR1 and Fas. A novel cell death receptor was recently identified by several groups independently and designated DR3, Wsl-1, Apo-3, TRAMP and LARD1-5. The ligand for this novel death receptor has been defined as TWEAK, also termed Apo3L. DR3 is highly expressed in the tissues enriched in lymphocytes including PBL, thymus and spleen. Like TNFR1, DR3 induces apoptosis and NF-κB activation.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.