

# Histone H2A.X(Phospho-Ser139) Conjugated Antibody

Catalog No: #C11268

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)
 Package Size: #C11268-AF350 100ul #C11268-AF405 100ul #C11268-AF488 100ul #C11268-AF555 100ul #C11268-AF594 100ul #C11268-AF647 100ul #C11268-AF680 100ul #C11268-AF750 100ul #C11268-Biotin 100ul
   
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Histone H2A.X(Phospho-Ser139) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of Histone H2A.X only when phosphorylated at serine 139.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 139 (Q-A-S(p)-Q-E) derived from Human Histone H2A.X.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Target Name	Histone H2A.X
Modification	Phospho-Ser139
Other Names	H2A.X; H2AFX; H2a/x; HIST5-2AX;
Accession No.	Swiss-Prot: P16104 NCBI Protein: NP_002096.1
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

## Application Details

### Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.

Yaneva M, et al. (2005) *Nucleic Acids Res.* 33(16): 5320-5330.

Tsukuda T, et al.(2006) *Nature*. Author manuscript; available in PMC 2006 March 6.

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.