

CIDE-B Antibody


Catalog No: #24116

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	CIDE-B Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB IHC
Species Reactivity	Hu Ms Rt
Specificity	CIDE-B antibody has no cross activity to CIDE-A.
Immunogen Type	Peptide
Immunogen Description	Raised against a peptide corresponding to 15 amino acids near the carboxy terminus of human CIDE-B.
Target Name	CIDE-B
Other Names	Cell death-inducing DFF-like effector B
Accession No.	AAF27658
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of CIDE-B in mouse small intestine tissue lysate with CIDE-B antibody at (A) 0.5, (B) 1 and (C) 2 ug/mL.



Immunohistochemistry of CIDE-B in human small intestine tissue with CIDE-B antibody at 5 ug/mL.

Background

Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. DFF45/ICAD has been identified as inhibitor of caspase activated DNase DFF40/CAD. DFF45 related proteins CIDE-A and CIDE-B (for cell death-inducing DFF-like effector A and B) were recently identified. CIDE contains a new type of domain termed CIDE-N, which has high homology with the regulatory domains of DFF45/ICAD and DFF40/CAD. Expression of CIDE-B induces apoptosis, which is inhibited by DFF45. CIDE-B is a DFF45-inhibitable effector that promotes cell death and DNA fragmentation. CIDE-B is expressed mainly in liver and at lower levels in spleen, kidney, peripheral blood lymphocytes and bone marrow.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.