Akt1 (phospho Ser246) Polyclonal Antibody

Catalog No: #14104

Description

Applications

Specificity

Species Reactivity

Package Size: #14104-1 50ul #14104-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	Akt1 (phospho Ser246) Polyclonal Antibody
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific
	immunogen.

	phosphorylated at 62 fe.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human Akt around the phosphorylation

Phospho-Akt1 (S246) Polyclonal Antibody detects endogenous levels of Akt1 protein only when

site of Ser246. AA range:212-261

WB,IHC-p,IF/ICC,ELISA

phosphorylated at \$246

Human, Mouse, Rat

Other Names AKT1; PKB; RAC; RAC-alpha serine/threonine-protein kinase; Protein kinase B; PKB; Protein kinase B alpha;

PKB alpha; Proto-oncogene c-Akt; RAC-PK-alpha

Accession No. Swiss Prot:P31749GeneID:207
SDS-PAGE MW 56

Concentration 1 mg/ml

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage -20°C/1

Application Details

Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

Background

AKT serine/threonine kinase 1(AKT1) Homo sapiens The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2011]

Note: This product is for in vitro research use only and is not intended for use in humans or animals.