McI-1 Antibody

Catalog No: #24300




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$		4.0
1	Decri	ption
\boldsymbol{L}	COUL	บแบบ

Product Name	McI-1 Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Ion exchange chromatography purified	
Applications	ELISA WB ICC	
Species Reactivity	Hu	
Specificity	Detects both isoforms of McI-1	
Immunogen Type	Peptide	
Immunogen Description	Raised against a peptide corresponding to 16 amino acids near the center of human Mcl-1.	
Target Name	McI-1	
Other Names	Myeloid cell leukemia 1	
Accession No.	NP_068779	
Concentration	1mg/ml	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

Images



Western blot analysis of Mcl-1 in Raji cell lysates with Mcl-1 antibody (IN) at (A) 0.5, (B) 1, and (C) 2 ug/mL.



Immunocytochemistry of McI-1 in Raji cells with McI-1 antibody at 10 ug/mL.

Background

Myeloid cell leukimia-1 (Mcl-1) is a member of the Bcl-2 family of proteins that can act to promote cell survival. While the mechanism by which Mcl-1 inhibits apoptosis is not known, it is thought that it may heterodimerize and neutralize pro-apoptotic members of the Bcl-2 family such as Bim or Bak. Mcl-1 was originally identified in differentiating myeloid cells, but has since been shown to be expressed in multiple cell types. Mcl-1 is essential for embryogenesis and for the development and maintenance of B and T lymphocytes in animals. Mcl-1 exists as at least two distinct isoforms designated Mcl-1L and Mcl-1S. In marked contrast to the larger isoform of Mcl-1, overexpression of Mcl-1S promotes cell death.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.