V5-tag Rabbit mAb

Catalog No: #56317

Package Size: #56317-1 50ul #56317-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | V5-tag Rabbit mAb |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB |
| Species Reactivity | All |
| Specificity | V5 tag antibody detects endogenous levels of C-terminal, internal, and N-terminal V5-tagged proteins. |
| Immunogen Description | A synthesized peptide |
| Other Names | V5 tag |
| Accession No. | Uniprot:N/A |
| Calculated MW | N/A |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Application Details

WB:1:3000~1:10000

Images



Western blot analysis of extracts from v5-tag fusion protein, using V5 tag antibody.

Product Description

The V5 epitope tag is derived from a small epitope (Pk) present on the P and V proteins of the paramyxovirus of simian virus 5 (SV5). The V5 tag is usually used with all 14 amino acids (GKPIPNPLLGLDST), although it has also been used with a shorter 9 amino acid sequence (IPNPLLGLD).

| Note: This product is for in vitro research use only and is not intended for use in humans or animals. | | | |
|--|--|--|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |