Chk2 (Phospho-Thr68) Conjugated Antibody

Catalog No: #C14265



Package Size: #C14265-AF350 100ul #C14265-AF405 100ul #C14265-AF488 100ul #C14265-AF555 100ul #C14265-AF55

| Description | |
|-----------------------|---|
| Product Name | Chk2 (Phospho-Thr68) Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB, IF |
| Species Reactivity | Human |
| Specificity | Phospho-Chk2 (T68) Antibody detects endogenous levels of total Phospho-Chk2 (T68) |
| Immunogen Description | A synthesized peptide derived from human Phospho-Chk2 (T68) |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | CHEK2; CHK2; Cds1; Chk2; EC 2.7.11.1; RAD53; kinase Chk2; |
| Accession No. | Uniprot:O96017 |
| Calculated MW | 65kDa |
| Storage | Store at 4°C in dark for 6 months |

Application Details

WB: 1:50-1:200 IF:1:50-1:200

Product Description

These are known to be preferred sites for phosphorylation by ATM/ATR kinases. After DNA damage by ionizing radiation (IR), UV irradiation, or hydroxyurea treatment, Thr68 and other sites in this region become phosphorylated by ATM/ATR. The SQ/TQ cluster domain, therefore, seems to have a regulatory function.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.