

Recombinant Human 4-1BB Receptor/TNFRSF9

Catalog No: #AP60042


Package Size: #AP60042-1 5ug #AP60042-2 100ug #AP60042-3 500ug

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Recombinant Human 4-1BB Receptor/TNFRSF9
Host Species	Escherichia coli.
Purification	> 97 % by SDS-PAGE and HPLC analyses.
Other Names	TNFRSF9, CD137 Antigen, T-cell Antigen ILA
Calculated MW	Approximately 17.7 kDa, a single non-glycosylated polypeptide chain containing 166 amino acids.
Target Sequence	ERTRSLQDPC SNCPAGTFCD NNRNQICSPC PPNSFSSAGG QRTCDICRQC KGVFRTRKEC SSTSNAECDG TPGFHCLGAG CSMCEQDCKQ GQELTKKGCK DCCFGTFNDQ KRGICRPWTN CSLDGKSVLV NGTKERDVVC GPSPADLSPG ASSVTPPAPA REPGHS
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 10 mM PB, pH 8.0, 150 mM NaCl.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.- 12 months from date of receipt, -20 to -70 °C as supplied.- 1 month, 2 to 8 °C under sterile conditions after reconstitution.- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

Images



Background

4-1BB receptor, also named TNFRSF9 is a member of the TNF superfamily of receptors. It is mainly expressed on the surface of a variety of T cells, but also found in B cells, monocytes, and various transformed cell lines. 4-1BB receptor binds to 4-1BBL, and they co-stimulate activity for activated T cells. Signaling by 4-1BB Receptor has been implicated in the antigen-presentation process and generation of cytotoxic T cells. Crosslinking of 4-1BB Receptor enhances T cell proliferation, IL-2 secretion survival and cytolytic activity. Further, it can enhance immune activity to eliminate tumors in mice.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.