

Sumo Antibody


Catalog No: #24467

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	Sumo Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB IHC
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide
Immunogen Description	Raised against a 14 amino acid peptide from near the amino terminus of human sumo.
Target Name	Sumo
Other Names	small-ubiquitin-like modifier 1, ubiquitin-like protein 1, GAP-modifying protein 1
Accession No.	AAH66306
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of sumo in HL-60 cell lysate with sumo antibody at (A) 0.5, (B) 1, and (C) 2 ug/mL.



Immunocytochemistry of Sumo in HL60 cells with Sumo antibody at 5 ug/mL.

Background

The sumo family of proteins is related both structurally and functionally to ubiquitin in that they are post-translationally attached to the e-amino group of a lysine residue of the substrate protein. This sumoylation plays a number of roles in DNA replication and repair, protein targeting to various subnuclear structures, and the regulation of numerous cellular processes including the inflammatory response in mammalian cells. Sumo was initially identified as a covalent modification of RanGAP1 in studies on nuclear import in mammalian cells. More recently, sumo has been shown to be involved in the regulation of transcription factors, possibly by enhancing their interactions with co-repressors. Sumo is also thought to play some role in the modulation of ubiquitin-mediated degradation of proteins by acting as an inhibitor. At least four different isoforms of sumo are known to exist; Sumo antibody will only recognize isoform 1.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.