## ATG12 Antibody

Catalog No: #24617




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


	es	cr	nt	חוי	n
-	-	UI	ועו	IIU.	

Product Name	ATG12 Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Affinity chromatography purified via peptide column	
Applications	ELISA WB IHC	
Species Reactivity	Hu Ms Rt	
Immunogen Type	Peptide	
Immunogen Description	Raised against a 16 amino acid peptide from near the amino terminus of human ATG12.	
Target Name	ATG12	
Other Names	Autophagy protein 12, Autophagy related protein 12, APG12, APG12L, HAPG12	
Accession No.	EAW48955	
Concentration	1mg/ml	
Formulation	Supplied in PBS containing 0.02% sodium azide.	
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated	
	freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	

## **Images**



Western blot analysis of ATG12 in human brain tissue lysate with ATG12 antibody at (A) 0.5, and (B) 1 ug/mL.



Immunohistochemistry of ATG12 in human brain tissue with ATG12 antibody at 2.5  $\,$  ug/mL.

## Background

Autophagy, the process of bulk degradation of cellular proteins through an autophagosomic-lysosomal pathway is important for normal growth control and may be defective in tumor cells. It is involved in the preservation of cellular nutrients under starvation conditions as well as the normal turnover of cytosolic components. This process is negatively regulated by TOR (Target of rapamycin) through phosphorylation of autophagy protein APG1. ATG12, another member of the autophagy protein family, forms a conjugate with ATG5; this conjugate has a ubiquitin-protein ligase (E3)-like activity for protein lipidation in autophagy. This conjugate also associates with innate immune response proteins such as RIG-I and VISA (also known as IPS-1), inhibiting type I interferon production and permitting viral replication in host cells. ATG12 has also been shown to interact with ATG10 in human embryonic kidney cells in the presence of ATG7. At least two isoforms of ATG12 are known to exist.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.