ATG5 Antibody

Catalog No: #24623




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| $\overline{}$ | | 4.0 |
|------------------|-------|-------|
| 1 | Decri | ption |
| \boldsymbol{L} | COUL | บแบบ |

| Product Name | ATG5 Antibody | |
|-----------------------|--|--|
| Host Species | Rabbit | |
| Clonality | Polyclonal | |
| Purification | Affinity chromatography purified via peptide column | |
| Applications | ELISA WB IHC | |
| Species Reactivity | Hu Ms Rt | |
| Specificity | Three isoforms of ATG5 are known to exist; this ATG5 antibody will only detect the longest isoform. | |
| Immunogen Type | Peptide | |
| Immunogen Description | Raised against a 16 amino acid peptide from near the amino terminus of human ATG5. | |
| Target Name | ATG5 | |
| Other Names | Autophagy protein 5, Autophagy related protein 5, ATG5L, ASP | |
| Accession No. | EAW48415 | |
| Concentration | 1mg/ml | |
| Formulation | Supplied in PBS containing 0.02% sodium azide. | |
| Storage | Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated | |
| | freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures. | |

Images



Western blot analysis of ATG5 in rat spleen tissue lysate with ATG5 antibody at (A) 1 and (B) 2 μ .



Immunohistochemistry of ATG5 in human spleen tissue with ATG5 antibody at 2.5 μ mL.

Background

Autophagy, the process of bulk degradation of cellular proteins through an autophagosomic-lysosomal pathway is important for normal growth control and may be defective in tumor cells. It is involved in the preservation of cellular nutrients under starvation conditions as well as the normal turnover of cytosolic components. This process is negatively regulated by TOR (Target of rapamycin) through phosphorylation of autophagy protein APG1. ATG5, another member of the autophagy protein family, forms a conjugate with ATG12; this conjugate has a ubiquitin-protein ligase (E3)-like activity for protein lipidation in autophagy. This conjugate also associates with innate immune response proteins such as RIG-I and VISA (also known as IPS-1), inhibiting type I interferon production and permitting viral replication in host cells.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.