## JPH3 Antibody

Catalog No: #24795




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _                |       |     |      |
|------------------|-------|-----|------|
|                  | escri | ınt | ion. |
| $\boldsymbol{L}$ | COUL  | Iυι | IUII |

| Product Name          | JPH3 Antibody  |  |
|-----------------------|--|--|
| Host Species          | Rabbit   |  |
| Clonality             | Polyclonal   |  |
| Purification          | Affinity chromatography purified via peptide column  |  |
| Applications          | ELISA WB IHC   |  |
| Species Reactivity    | Hu Ms Rt   |  |
| Immunogen Type        | Peptide  |  |
| Immunogen Description | Raised against a 18 amino acid peptide near the carboxy terminus of human JPH3.                            |  |
| Target Name           | JPH3   |  |
| Other Names           | Junctophilin 3, JP3, JP-3, TNRC22, HDL2  |  |
| Accession No.         | NP_065706  |  |
| Concentration         | 1mg/ml   |  |
| Formulation           | Supplied in PBS containing 0.02% sodium azide.   |  |
| Storage               | Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated |  |
|                       | freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.                       |  |

## **Images**



Western blot analysis of JPH3 in Daudi cell lysate with JPH3 antibody at 1 ug/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of JPH3 in human brain tissue with JPH3 antibody at 2.5  $\mu$ ML.

## Background

Junctional complexes between the plasma membrane (PM) and endoplasmic/sarcoplasmic reticulum (ER/SR) are a common feature of all excitable cell types and mediate cross talk between cell surface and intracellular ion channels. Junctophilins (JPs) are important components of the junctional complexes. JPs are composed of a carboxy-terminal hydrophobic segment spanning the ER/SR membrane and a remaining cytoplasmic domain that shows specific affinity for the PM. Four JPs have been identified as tissue-specific subtypes derived from different genes: JPH1 is expressed in skeletal muscle, JPH2 is detected throughout all muscle cell types, and JPH3 and JPH4 are predominantly expressed in the brain. In the CNS, both JPH3 and JPH4 are expressed throughout neural sites and contribute to the subsurface cistern formation in neurons. Mice lacking both JPH3 and JPH4 subtypes exhibit serious symptoms such as impaired learning and memory and are accompanied by abnormal nervous functions. A repeat expansion in JPH3 is associated with Huntington disease-like 2. At least two isoforms of JPH3 are known to exist.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.