Wee1(Phospho-Ser123) Antibody Biotin Conjugated

Catalog No: #C05615B



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | Support: tech@signalwayantibody.com |
|-----------------------|---|
| Product Name | Wee1(Phospho-Ser123) Antibody Biotin Conjugated |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | lgG |
| Purification | Purified by Protein A. |
| Applications | WB IHC-P |
| Species Reactivity | Hu Ms Rt |
| Immunogen Description | KLH conjugated synthetic phosphopeptide aa 100-150 646 derived from human Wee1 around the |
| | phosphorylation site of Ser123 [SS(p-S)PV] |
| Conjugates | Biotin |
| Target Name | Wee1 Ser123 |
| Other Names | WEE1A; WEE1hu; Wee1-like protein kinase; Wee1A kinase; WEE1 |
| Accession No. | Swiss-Prot#P30291NCBI Gene ID7465 |
| Cell Localization | Nucleus |
| Concentration | 1mg ml |
| Formulation | 10mM Tris Buffered Saline containing 1% BSA, 50% glycerol and 0.09% sodium azide. |
| Storage | Store at 4C for 12 months. |

Application Details

Western blotting: 1:100-1000lmmunohistochemistry1:100-500

Background

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase. Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M G1 phase, probably due to its degradation.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.