## DR5 Monoclonal Antibody

Catalog No: #27200

Package Size: #27200-1 50ul #27200-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


| Beeenplien            |  |
|-----------------------|--|
| Product Name          | DR5 Monoclonal Antibody  |
| Host Species          | Mouse  |
| Clonality             | Monoclonal   |
| Clone No.             | 7F4-F8-G11   |
| Isotype               | lgG1   |
| Purification          | Affinity purified  |
| Applications          | WB ICC   |
| Species Reactivity    | Hu Ms  |
| Specificity           | This antibody detects endogenous levels of TNFRSF10B and does not cross-react with related proteins.   |
| Immunogen Type        | Recombinant Protein  |
| Immunogen Description | Purified recombinant human TNFRSF10B protein fragments expressed in E.coli.                            |
| Target Name           | DR5  |
| Other Names           | Fas like protein; Apoptosis inducing protein TRICK2A/2B; Apoptosis inducing receptor TRAIL R2; CD 262; |
|                       | CD262; CD262 antigen; Cytotoxic TRAIL receptor 2; Death domain containing receptor for TRAIL/Apo 2L;   |
|                       | Death domain containing receptor for TRAIL/Apo2L;  |
| Accession No.         | Uniprot: O14763 Gene ID: 8795  |
| SDS-PAGE MW           | 48kd   |
| Formulation           | Purified mouse monoclonal in PBS(pH 7.4) containing with 0.02% sodium azide,0.1mg/mlBSA and 50%        |
|                       | glycerol.  |
| Storage               | store at -20A C  |
|                       |  |

## Application Details Western blotting: 1:500-1:2000 Immunocytochemistry: 1:100

## Images



Immunocytochemistry of HeLa cells fixed by Paraformaldehyde and using anti-TNFRSF10B antibody diluted 1:100.



Western blot detection of TNFRSF10B antibody in Mouse kindey, Mouse brain, HCT116 and Hela cell lysates using TNFRSF10B antibody (1:500-1:2000 diluted). Predicted band size: 40/48KDa. Observed band size: 40/48KDa.

## Background

Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B.Essential for ER stress-induced apoptosis.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.