

## IRF-3 (Phospho-Ser386) Antibody

Catalog No: #11760



Package Size: #11760-1 50ul #11760-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	IRF-3 (Phospho-Ser386) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of IRF-3 only when phosphorylated at serine 386.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of Serine 386(A-S-S(p)-L-E) derived from Human IRF-3.
Target Name	IRF-3
Modification	Phospho
Other Names	IRF3; Interferon regulatory factor 3;
Accession No.	Swiss-Prot#: Q14653; NCBI Gene#: 3661; NCBI Protein#: NP_001184052.1.
SDS-PAGE MW	55kd
Concentration	0.8mg/ml
Formulation	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C/1 year

## Application Details

Western blotting: 1:500~1:1000

## Images



Western blot analysis of extracts from untreated and treated 293T cell lysate using IRF-3 (Phospho-Ser386) Antibody #11760 at 1/500 dilution.

## Background

---

Key transcriptional regulator of type I interferon (IFN)-dependent immune responses and plays a critical role in the innate immune response against DNA and RNA viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters. Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha (IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene induction. Found in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, becomes phosphorylated by IKKBE and TBK1 kinases. This induces a conformational change, leading to its dimerization and nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes. Can activate distinct gene expression programs in macrophages and can induce significant apoptosis in primary macrophages.

Au W.W.-C., Proc. Natl. Acad. Sci. U.S.A. 92:11657-11661(1995).

The MGC Project Team; Genome Res. 14:2121-2127(2004).

Bellingham J., Ann. Hum. Genet. 62:231-234(1998).

---

Note: This product is for in vitro research use only and is not intended for use in humans or animals.