CAD (Phospho-Thr456) Antibody

Catalog No: #11789

Package Size: #11789-1 50ul #11789-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | |
|---|-------|-----|----|
| | escri | ntı | Λn |
| u | COUL | υu | UH |
| | | | |

| Product Name | CAD (Phospho-Thr456) Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | IHC |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of CAD only when phosphorylated at threonine 456. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of threonine 456 (P-I-T(p)-P-H) derived from Human CAD. |
| Target Name | CAD |
| Modification | Phospho |
| Other Names | PYR1; CAD protein; EC 2.1.3.2; EC 3.5.2.3; |
| Accession No. | Swiss-Prot#: P27708; NCBI Gene#: 790; NCBI Protein#: NP_004332.2. |
| SDS-PAGE MW | 242kd |
| Concentration | 1.0mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide |
| | and 50% glycerol. |
| Storage | Store at -20°C/1 year |
| | |

Application Details

Immunohistochemistry: 1:50~1:100

Images



Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue using CAD (Phospho-Thr456) antibody #11789 (left)or the same antibody preincubated with blocking peptide (right).

Background

The de novo synthesis of pyrimidine nucleotides is required for mammalian cells to proliferate. This gene encodes a trifunctional protein which is associated with the enzymatic activities of the first 3 enzymes in the 6-step pathway of pyrimidine biosynthesis: carbamoylphosphate synthetase (CPS II), aspartate transcarbamoylase, and dihydroorotase. This protein is regulated by the mitogen-activated protein kinase (MAPK) cascade, which indicates a direct link between activation of the MAPK cascade and de novo biosynthesis of pyrimidine nucleotides.

Iwahana H., Biochem. Biophys. Res. Commun. 219:249-255(1996).

Davidson J.N., DNA Cell Biol. 9:667-676(1990).

Olsen J.V., Cell 127:635-648(2006).

Note: This product is for in vitro research use only and is not intended for use in humans or animals.