## CRMP-2 (Phospho-Thr509) Antibody

Catalog No: #11795

Package Size: #11795-1 50ul #11795-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description			
Product Name	CRMP-2 (Phospho-Thr509) Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.		
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho		
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.		
Applications	WB IHC		
Species Reactivity	Hu		
Specificity	The antibody detects endogenous levels of CRMP-2 only when phosphorylated at threonine 509.		
Immunogen Type	Peptide-KLH		
Immunogen Description	Peptide sequence around phosphorylation site of threonine 509 (S-V-T(p)-P-K) derived from Human CRMP-2.		
Target Name	CRMP-2		
Modification	Phospho		
Other Names	CRMP2; N2A3; TOAD-64; DPYL2; ULIP2		
Accession No.	Swiss-Prot#: Q16555; NCBI Gene#: 1808; NCBI Protein#: NP_001377.1.		
SDS-PAGE MW	50kd		
Concentration	1.0mg/ml		
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCI, 0.02% sodium azide		
	and 50% glycerol.		

pplication Details	etails	ation Details
/estern blotting: 1:500~1:100	:500~1:1000	blotting: 1:500~1:100
nmunohistochemistry: 1:50~	istry: 1:50~1:1	nistochemistry: 1:50~1


Store at -20°C/1 year

## Images

Storage

	HT29 HT29				
			117		
			85		
	CRMP-2		10		
	(pThr509)		48		
	, , , , , , , , , , , , , , , , , , ,		34		
			26		
			19		
			(kD)		

Western blot analysis of extracts from HT-29 cells treated with heat shock using CRMP-2 (Phospho-Thr509) Antibody #11795.The lane on the right is treated with the antigen-specific peptide.



Immunohistochemical analysis of paraffin-embedded human brain tissue using CRMP-2 (Phospho-Thr509) antibody #11795 (left)or the same antibody preincubated with blocking peptide (right).

## Background

CRMP-2 is an enzyme with dihydropyrimidinase activity. Plays a role in RhoA-dependent signaling, through interaction with and regulation of Rho kinase. Plays a role in neurogenesis. Aberrantly expressed in fetal Down syndrome brain. Miki Y., Science 266:66-71(1994). Smith T.M., Genome Res. 6:1029-1049(1996).

Wilson C.A., Oncogene 14:1-16(1997).

Note: This product is for in vitro research use only and is not intended for use in humans or animals.