YB1 (Phospho-Ser102) Antibody

Catalog No: #11819

Package Size: #11819-1 50ul #11819-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | | | |
|-----------------------|---|--|--|
| Product Name | YB1 (Phospho-Ser102) Antibody | | |
| Host Species | Rabbit | | |
| Clonality | Polyclonal | | |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. | | |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho | | |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. | | |
| Applications | WB | | |
| Species Reactivity | Hu Ms Rt | | |
| Specificity | The antibody detects endogenous levels of YB1 only when phosphorylated at serine 102. | | |
| Immunogen Type | Peptide-KLH | | |
| Immunogen Description | Peptide sequence around phosphorylation site of Serine 102(L-R-S(p)-V-G) derived from Human YB1. | | |
| Target Name | YB1 | | |
| Modification | Phospho | | |
| Other Names | CBF-A; NSEP1; EFI-A; MSY-1; YBX1 | | |
| Accession No. | Swiss-Prot#: P67809; NCBI Gene#: 4904; NCBI Protein#: NP_004550.2. | | |
| SDS-PAGE MW | 36kd | | |
| Concentration | 1.0mg/ml | | |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide | | |
| | and 50% glycerol. | | |
| Storage | Store at -20°C/1 year | | |

Application Details

Western blotting: 1:500~1:1000

Images

| | HepG2 HepG2 | | |
|---------------------------------------|---------------|------|--|
| | | 117 | |
| | | 85 | |
| | | | |
| | | 48 | |
| (pSe | YB1 er102) | 34 | |
| , , , , , , , , , , , , , , , , , , , | | 26 | |
| | | 19 | |
| | | (kD) | |
| | | | |

Western blot analysis of extracts from HepG2 cells treated with PMA using YB1 (Phospho-Ser102) Antibody #11819.The lane on the right is treated with the antigen-specific peptide.

Background

YB-1 is a nuclear protein that binds to splice sites in pre-mRNA and regulates splice site selection. Binds and stabilizes cytoplasmic mRNA. Contributes to the regulation of translation by modulating the interaction between the mRNA and eukaryotic initiation factors CCAAT-containing Y-box of HLA class II genes. Component of cytoplasmic messenger ribonucleoprotein particles (mRNPs). Interacts with AKT1, SFRS9, THOC4, MSH2, XRCC5, WRN and NCL. Can bind to DNA as a homomeric form, (EFI-A)n or as a heteromeric form in association with EFI-B. Homodimer in the presence of ATP.

Sakura H., Gene 73:499-507(1988).

Didier D.K. Proc. Natl. Acad. Sci. U.S.A. 85:7322-7326(1988).

Kolluri R., Nucleic Acids Res. 19:4771-4771(1991).

Note: This product is for in vitro research use only and is not intended for use in humans or animals.