NFKBIA Antibody

Catalog No: #32213

Package Size: #32213-1 50ul #32213-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| | 1. T. C. |
|-----------------------|--|
| Product Name | NFKBIA Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were purified by affinity purification using immunogen. |
| Applications | WB |
| Species Reactivity | Human,Mouse,Rat |
| Specificity | The antibody detects endogenous level of total NFKBIA protein. |
| Immunogen Type | Recombinant Protein |
| Immunogen Description | Recombinant protein of human NFKBIA. |
| Target Name | NFKBIA |
| Other Names | NFKBIA; IKBA; MAD-3; NFKBI; |
| Accession No. | Swiss-Prot:P25963NCBI Gene ID:4792 |
| SDS-PAGE MW | 39KD |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C |
| | |

Application Details

WB 1:500 - 1:2000

Images



Western blot analysis of extracts of various cell lines, using IKB alpha antibody at 1:1000 dilution.

Background

The NF-κB/Rel transcription factors are present in the cytosol in an inactive state complexed with the inhibitory IκB proteins (1-3). Activation occurs via phosphorylation of IκBα at Ser32 and Ser36 followed by proteasome-mediated degradation that results in the release and nuclear translocation of

active NF-κB (3-7). IκBα phosphorylation and resulting Rel-dependent transcription are activated by a highly diverse group of extracellular signals including inflammatory cytokines, growth factors, and chemokines. Kinases that phosphorylate IκB at these activating sites have been identified (8).

Note: This product is for in vitro research use only and is not intended for use in humans or animals.