

QK1 Polyclonal Antibody Cy5.5 Conjugated

Catalog No: #C01318Cy5.5

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	QK1 Polyclonal Antibody Cy5.5 Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	IF(IHC-P)
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human QK1
Conjugates	Cy5.5
Target Name	QK1
Other Names	HKQ; Homolog of mouse quaking QKI KH domain RNA binding protein; Hqk; HQK1; Hqkl; Protein quaking; QK; QK1; QK3; QKI; QKI_HUMAN; QKI1; Quaking homolog; Quaking homolog KH domain RNA binding; Quaking homolog KH domain RNA binding mouse; Quaking isoform 1; Quaking protein; RNA binding protein HQK.
Concentration	1mg ml
Formulation	Aqueous buffered solution containing 1% BSA, 50% glycerol and 0.09% sodium azide.
Storage	Store at 4C for 12 months.

Application Details

IF:1:50-200

Background

QKI is a 341 amino acid protein that localizes to both the cytoplasm and the nucleus and contains one KH domain. Expressed in the frontal cortex of the brain, QKI functions as an RNA-binding protein that plays an important role in myelination and specifically binds to the RNA core sequence 5'-NACUAAY-N(1,20)-UAAY-3'. Additionally, QKI regulates pre-mRNA splicing, and mRNA export and is involved in protecting and promoting the stability of select mRNAs. QKI may be methylated by PRMT1 and may also be phosphorylated at its C-terminus, an event that decreases QKI mRNA-binding affinity. Defects or deletions in the gene encoding QKI are associated with astrocytic tumors and may be involved in the pathogenesis of schizophrenia. Multiple isoforms of QKI exist due to alternative splicing events.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.