Description

## JIP3 Polyclonal Antibody Cy5.5 Conjugated

Catalog No: #C02749Cy5.5



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description           |  |
|-----------------------|--|
| Product Name          | JIP3 Polyclonal Antibody Cy5.5 Conjugated  |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Isotype               | IgG  |
| Purification          | Purified by Protein A.   |
| Applications          | IF(IHC-P)  |
| Species Reactivity    | Hu Ms Rt   |
| Immunogen Description | KLH conjugated synthetic peptide derived from human JIP3   |
| Conjugates            | Cy5.5  |
| Target Name           | JIP3   |
| Other Names           | C-jun-amino-terminal kinase-interacting protein 3; FLJ00027; Homolog of Drosophila Sunday driver 2; JIP-3; |
|                       | JIP3_HUMAN; JNK Stress Activated Protein Kinase Associated Protein 1; JNK Interacting Protein 3; JNK       |
|                       | MAP kinase scaffold protein 3; JNK-interacting protein 3; KIAA1066; MAPK8IP3; Mitogen Acti                 |
| Concentration         | 1mg ml   |
| Formulation           | Aqueous buffered solution containing 1% BSA, 50% glycerol and 0.09% sodium azide.                          |
| Storage               | Store at 4C for 12 months.   |

## **Application Details**

IF:1:50-200

## Background

The JNK-interacting proteins (JIPs) are a family of scaffold proteins that mediate JNK signaling by organizing specific components of the MAPK cascade together to form a functional JNK signaling molecule. JIP-3 (JNK-interacting protein 3), also known as JSAP1 or MAPK8IP3 (Mitogen-activated protein kinase 8-interacting protein 3), is a 1,336 amino acid protein that localizes to the cytoplasm and belongs to the JIP family. Expressed in a variety of tissues, including brain and heart, JIP-3 forms homo- or heterooligomeric complexes that can interact with several components of the JNK signaling pathway, thereby functioning as a regulator of kinesin-dependent axonal transport that may also play a role in scaffold formation within neuronal cells. Human JIP-3, which may be phosphorylated upon DNA damage, shares 69% similarity with its mouse counterpart, suggesting a conserved role between species. Multiple isoforms of JIP-3 exist due to alternative splicing events.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.