

COT antibody

Catalog No: #22076

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	COT antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC
Species Reactivity	Hu
Immunogen Type	Recombinant protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 140 and 432 of COT
Target Name	COT
Other Names	COT
Accession No.	NCBI Gene ID: 54677NCBI mRNA#: NM_021151NCBI Protein#: NP_066974
Concentration	1mg/ml
Formulation	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 70kd

Western blotting: 1:500-1:3000

Immunohistochemistry: 1:100-1:250

Images




Sample (30 ug of whole cell lysate)

A: 293T

B: A431

7.5% SDS PAGE

COT antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded Ca922 xenograft, using CROT antibody at 1: 100 dilution.

Background

This gene encodes a member of the carnitine/choline acetyltransferase family. The encoded protein converts 4,8-dimethylnonanoyl-CoA to its corresponding carnitine ester. This transesterification occurs in the peroxisome and is necessary for transport of medium- and long- chain acyl-CoA molecules out of the peroxisome to the cytosol and mitochondria. The protein thus plays a role in lipid metabolism and fatty acid beta-oxidation. Alternatively spliced transcript variants have been described.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.