p95/NBS1(Phospho-Ser343) Antibody

Catalog No: #11057

Package Size: #11057-1 50ul #11057-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	p95/NBS1(Phospho-Ser343) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB
Species Reactivity	Ни
Specificity	The antibody detects endogenous level of p95/NBS1 only when phosphorylated at serine 343.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 343 (S-L-S(p)-Q-G) derived from Human p95/NBS1.
Target Name	p95/NBS1
Modification	Phospho
Other Names	NBN
Accession No.	Swiss-Prot: O60934NCBI Protein: NP_002476.2
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 95kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from Hela cells untreated or treated with Hydroxyurea using p95/NBS1(Phospho-Ser343) Antibody #11057.

Background

Mutations in p95/NBS1 gene are associated with Nijmegen breakage syndrome, an autosomal recessive chromosomal instability syndrome characterized by microcephaly, growth retardation, immunodeficiency, and cancer predisposition. The encoded protein is a member of the MRE11/RAD50 double-strand break repair complex which consists of 5 proteins. This gene product is thought to be involved in DNA double-strand break repair and DNA damage-induced checkpoint activation. Hsu HL, et al (2005) Oncogene; 24(31): 4956-64. Falck J, et al. (2005) Nature; 434(7033): 605-11.

Buscemi G, et al. (2004) Oncogene; 23(46): 7691-700.

Beausoleil SA, et al. (2004) Proc Natl Acad Sci U S A; 101(33): 12130-5.

Published Papers

el at., Inhibition of TPL2 by interferon-α suppresses bladder cancer through activation of PDE4D.In J Exp Clin Cancer Res. On 2018 Nov 27 by Qiang Z, Zhou ZY et al..PMID: 30482227, , (2018)

PMID:30482227

Rachid Drissi, Jing Wu, Yafang Hu el at., Telomere Shortening Alters the Kinetics of the DNA Damage

Response after Ionizing Radiation in Human Cells, Cancer Prevention Research, 4(12):1973n— C1981(2011)

PMID:21930799

el at., Telomere Shortening Alters the Kinetics of the DNA Damage Response After Ionizing Radiation in Human Cells. In Cancer Prev Res (Phila) on 2011 Dec by Rachid Drissi, Jing Wu, et al.. PMID:21930799, , (2011)

PMID:21930799

F Carrillo, SA Schneider, AMR Taylor el at., Prominent Oromandibular Dystonia and Pharyngeal Telangiectasia in Atypical Ataxia Telangiectasia.,

Cerebellum, 8:22n- C27(2009)

PMID:18846412

el at., Prominent oromandibular dystonia and pharyngeal telangiectasia in atypical ataxia telangiectasia. In Cerebellum on 2009 Mar by Carrillo

F,Schneider SA,et al..PMID:18846412, , (2009)

PMID:18846412

Note: This product is for in vitro research use only and is not intended for use in humans or animals.