

CHOP (Phospho-Ser30) Antibody

Catalog No: #11976



Package Size: #11976-1 50ul #11976-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	CHOP (Phospho-Ser30) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB
Species Reactivity	Human
Specificity	The antibody detects endogenous level of CHOP only when phosphorylated at serine 30.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine30(V-L-S(p)-S-D) derived from Human CHOP .
Conjugates	Unconjugated
Target Name	CHOP
Modification	Phospho
Other Names	C/EBP-homologous protein; DDIT3; GADD153; DNA-damage inducible transcript 3;
Accession No.	Swiss-Prot#: P35638; NCBI Gene#: 1649; NCBI Protein#: NP_001181982.1
SDS-PAGE MW	19kd
Concentration	1.0mg/ml
Formulation	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C/1 year

Application Details

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from Jurkat cells treated with PMA using Phospho-CHOP (Ser30) antibody #11976. The lane on the right is treated with the antigen-specific peptide.

Background

Multifunctional transcription factor in ER stress response. Plays an essential role in the response to a wide variety of cell stresses and induces cell cycle arrest and apoptosis in response to ER stress. Plays a dual role both as an inhibitor of CCAAT/enhancer-binding protein (C/EBP) function and as an activator of other genes. Acts as a dominant-negative regulator of C/EBP-induced transcription: dimerizes with members of the C/EBP family, impairs their association with C/EBP binding sites in the promoter regions, and inhibits the expression of C/EBP regulated genes. Positively regulates the transcription of TRIB3, IL6, IL8, IL23, TNFRSF10B/DR5, PPP1R15A/GADD34, BBC3/PUMA, BCL2L11/BIM and ERO1L.

Ubeda M, Habener JF (2003) J Biol Chem 278, 40514-20

Note: This product is for in vitro research use only and is not intended for use in humans or animals.