

Smad2(Phospho-S255) Rabbit mAb

Catalog No: #13429



Package Size: #13429-1 50ul #13429-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	Smad2(Phospho-S255) Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JF0882
Purification	ProA affinity purified
Applications	WB,IHC
Species Reactivity	Human;Mouse;Rat
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Ser255 of human Smad2.
Conjugates	Unconjugated
Other Names	Drosophila, homolog of, MADR2 antibody hMAD-2 antibody HsMAD2 antibody JV18 antibody JV18-1 antibody JV181 antibody MAD antibody MAD homolog 2 antibody MAD Related Protein 2 antibody Mad-related protein 2 antibody MADH2 antibody MADR2 antibody MGC22139 antibody MGC34440 antibody Mother against DPP homolog 2 antibody Mothers against decapentaplegic homolog 2 antibody Mothers against decapentaplegic, Drosophila, homolog of, 2 antibody Mothers against DPP homolog 2 antibody OTTHUMP00000163489 antibody Sma and Mad related protein 2 antibody Sma- and Mad-related protein 2 MAD antibody SMAD 2 antibody SMAD family member 2 antibody SMAD, mothers against DPP homolog 2 antibody SMAD2 antibody SMAD2_HUMAN antibody
Accession No.	Swiss-Prot#:Q15796
Calculated MW	52 kDa
SDS-PAGE MW	60 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:2000

IHC: 1:50-1:200

Images



All lanes: Smad2(Phospho-S255) Rabbit mAb at 1/1k dilution

Lane 1 : Hela whole cell lysates

Lane 2 : Hela treated with 200nM Calyculin A for 1 hours whole cell lysates

Lysates/proteins at 20 μ g per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution


Predicted band size: 52 kDa

Observed band size: 60 kDa

Exposure time: 15 seconds



Formalin-fixed;paraffin-embedded human uterus tissue stained for Smad2 (Phospho-S255) using 13429 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed;paraffin-embedded human bladder tissue stained for Smad2 (Phospho-S255) using 13429 at 1/100 dilution in immunohistochemical analysis.

Background

Smad proteins, the mammalian homologs of the *Drosophila* mothers against decapentaplegic (Mad), have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and Activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to Activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad proteins.

References

1. Ungefroren H et al. Rac1b negatively regulates TGF- β 1-induced cell motility in pancreatic ductal epithelial cells by suppressing Smad signalling. *Oncotarget* 5:277-90 (2014).
2. Harazono Y et al. miR-655 is an EMT-suppressive MicroRNA targeting ZEB1 and TGFBR2. *PLoS One* 8:e62757 (2013).

Published Papers

el., Gastrodin attenuates renal injury and collagen deposition via suppression of the TGF- β 1/Smad2/3 signaling pathway based on network

Note: This product is for in vitro research use only and is not intended for use in humans or animals.