

NF-κB p65 (Phospho-Ser536) Rabbit mAb

Catalog No: #14319



Package Size: #14319-1 50ul #14319-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	NF-κB p65 (Phospho-Ser536) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Purification	Protein A
Applications	WB;ICC/IF
Species Reactivity	Human;Mouse;Rat
Specificity	Phospho-NFκB-p65 (S536) Antibody detects endogenous levels of NFκB-p65 protein only when phosphorylated at S536.
Conjugates	Unconjugated
Target Name	NFκB p65
Modification	Phospho
Other Names	RELA;NFKB3;Transcription factor p65;Nuclear factor NF-κappa-B p65 subunit;Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3
Accession No.	Q04206
Calculated MW	60 kDa
SDS-PAGE MW	65 kDa
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Storage	Store at -20°C/1 year

Application Details



WB: 1:500-1:2000

ICC/IF: 1:50-1:200

Images



All lanes : NF-κB p65 (Phospho-Ser536) Rabbit mAb at 1/1k dilution
 Lane 1 : Hela whole cell lysates
 Lane 2 : Hela treated with 20ng/ml TNF-α for 15 minutes whole cell lysates
 Lane 3 : C2C12 whole cell lysates
 Lane 4 : C6 whole cell lysates
 Proteins at 20 μg per lane. Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 60 kDa Observed band size: 65 kDa
 Exposure time: 10 seconds



Immunocytochemistry/Immunofluorescence NF-κB p65 (Phospho Ser536) antibody (14319)
ICC/IF staining of NF-κB p65 (Phospho Ser536) in HeLa cells.
Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.
Samples were incubated with 14319 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 647 goat anti rabbit; used at a dilution of 1/500.
Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.