

CDC42 Rabbit mAb

Catalog No: #49282



Package Size: #49282-1 50ul #49282-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

| | |
|-----------------------|--|
| Product Name | CDC42 Rabbit mAb |
| Host Species | Recombinant Rabbit |
| Clonality | Monoclonal antibody |
| Clone No. | JJ086-04 |
| Purification | ProA affinity purified |
| Applications | WB, IHC, IP, FC |
| Species Reactivity | Human;Mouse;Rat |
| Immunogen Description | recombinant protein |
| Conjugates | Unconjugated |
| Other Names | CDC42 antibody CDC42_HUMAN antibody CDC42Hs antibody Cell division control protein 42 homolog antibody Cell division cycle 42 (GTP binding protein 25kDa) antibody Cell division cycle 42 antibody dJ224A6.1.1 (cell division cycle 42 (GTP-binding protein, 25kD)) antibody dJ224A6.1.2 (cell division cycle 42 (GTP-binding protein, 25kD)) antibody G25K antibody G25K GTP-binding protein antibody Growth regulating protein antibody GTP binding protein 25kDa antibody Small GTP binding protein CDC42 antibody TKS antibody |
| Accession No. | Swiss-Prot#:P60953 |
| Calculated MW | 21 kDa |
| SDS-PAGE MW | 21 kDa |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |


Application Details

WB: 1:1,000IHC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of CDC42 on different lysates using anti-CDC42 antibody at 1/1,000 dilution. Positive control: Lane 1: HeLa Lane 2: Jurkat Lane 3: HepG2




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-CDC42 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human pancreas tissue using anti-CDC42 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue using anti-CDC42 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of HeLa cells with CDC42 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

The superfamily of GTP-binding proteins, for which the Ras proteins are prototypes, has been implicated in regulation of diverse biological activities involving various aspects of cell growth and division. One mammalian member of the family, Cdc42, has an amino acid sequence that is similar to those of various members of the Ras superfamily proteins, including N-, K- and H-Ras, Rho proteins and the Rac proteins. On the basis of in vitro phosphorylation studies, it has been suggested that human Cdc42 may function in the signaling pathway of the EGF receptor or related growth factor receptor protein kinases. The Dbl oncogene has been shown to specifically catalyze dissociation of GDP from human Cdc42.

References

1. Gerasimcik N et al. The Rho GTPase Cdc42 Is Essential for the Activation and Function of Mature B Cells. J Immunol 194:4750-8 (2015). 2. Francis MK et al. Endocytic membrane turnover at the leading edge is driven by a transient interaction between Cdc42 and GRAF1. J Cell Sci 128:4183-95 (2015).

Published Papers

el at., A transcribed ultraconserved noncoding RNA, uc.285+, promotes colorectal cancer proliferation through dual targeting of CDC42 by directly binding mRNA and protein. In Transl Res on 2024 Aug by Sixian Chen, Qingyun Zhao,et al..PMID:38552953, , (2024)

[PMID:38552953](#)

Chen Sixian;Zhao Qingyun;Zhang Ruirui;Liu Jungang;Peng Wenyi;Xu Haotian;Li Xiaofei;Wang Xin;Wu Shuilian;Li Gang;Nan Aruo el at., A transcribed ultraconserved noncoding RNA, uc.285+, promotes colorectal cancer proliferation through dual targeting of CDC42 by directly binding mRNA and protein, , (2024)

PMID:

Note: This product is for in vitro research use only and is not intended for use in humans or animals.