

S100 beta Conjugated Antibody

Catalog No: #C48942



Package Size: #C48942-Biotin 100ul #C48942-AF350 100ul #C48942-AF405 100ul #C48942-AF488 100ul #C48942-AF555 100ul #C48942-AF594 100ul #C48942-AF647 100ul #C48942-AF680 100ul #C48942-AF750 100ul #C48942-Conjugated 50ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

Description

Product Name	S100 beta Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Applications	WB, IF
Species Reactivity	Hu, Ms, Rt, Goat, zebrafish
Immunogen Description	recombinant protein
Conjugates	AF0• §ε η´
Other Names	NEF antibody Protein S100 B antibody Protein S100-B antibody S 100 calcium binding protein beta chain antibody S 100 protein beta chain antibody S-100 protein beta chain antibody S-100 protein subunit beta antibody S100 antibody S100 calcium binding protein beta (neural) antibody S100 calcium-binding protein B antibody S100 protein beta chain antibody S100B antibody S100B_HUMAN antibody S100beta antibody
Accession No.	Swiss-Prot#:P04271
Calculated MW	11 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

WB: 1:50-1:200

IF:1:50-1:200

Background

The family of EF-hand type Ca2+-binding proteins includes calbindin (previously designated vitamin D-dependent Ca2+-binding protein), S-100 α and β, calgranulins A (also designated MRP8), B (also designated MRP14) and C (S-100 like proteins), and the parvalbumin family members, including parvalbumin α and parvalbuminβ(also designated oncomodulin). The S-100 protein is involved in the regulation of cellular processes such as cell cycle progression and differentiation. Research also indicates that the S-100 protein may function in the activation of Ca2+ induced Ca2+ release, inhibition of microtubule assembly and inhibition of protein kinase C mediated phosphorylation. Two S-100 subunits, sharing 60% sequence identity, have been described as S-100 α chain and S-100 β chain. Three S-100 dimeric forms have been characterized, differing in their subunit composition of either two α chains, two β chains or one α and one β chain. S-100 localizes to the cytoplasm and nuclei of astrocytes, Schwann's cells, ependymomas and astroglomas. S-100 is also detected in almost all benign naevi, malignant melanocytic tumours and in Langerhans cells in the skin. Calbindin, S-100 proteins and parvalbumin proteins are each expressed in neural tissues. In addition, S-100 α and β are present in a variety of other tissues, and calbindin is present in intestine and kidney.

Published Papers

el at., Honey Enriched with Additives Alleviates Behavioral, Oxidative Stress, and Brain Alterations Induced by Heavy Metals and Imidacloprid in Zebrafish. In Int J Mol Sci on 2024 Oct 31 by Emanuela Paduraru, Roxana Jijie,et al..PMID:39519279, , (2024)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.